

# Graphs and Coloring

Junaid

Lecture 7: Discrete Mathematical Modelling

## What is a Graph?

- ▶ A graph consists of two sets  $V$  and  $E$  where

## What is a Graph?

- ▶ A graph consists of two sets  $V$  and  $E$  where
- ▶

$$V = \{v_1, \dots, v_n\}$$

## What is a Graph?

- ▶ A graph consists of two sets  $V$  and  $E$  where

$$V = \{v_1, \dots, v_n\}$$

- ▶

$$E = \{e_1, \dots, e_m\}$$

## What is a Graph?

- ▶ A graph consists of two sets  $V$  and  $E$  where
- ▶
$$V = \{v_1, \dots, v_n\}$$
- ▶
$$E = \{e_1, \dots, e_m\}$$
- ▶ where each edge  $e_i$  consists of an **unordered** vertex pair  $e_i = \{v_j, v_k\}$ .

## What is a Graph?

- ▶ A graph consists of two sets  $V$  and  $E$  where
- ▶  
$$V = \{v_1, \dots, v_n\}$$
- ▶  
$$E = \{e_1, \dots, e_m\}$$
- ▶ where each edge  $e_i$  consists of an **unordered** vertex pair  $e_i = \{v_j, v_k\}$ .
- ▶  $j \neq k$  because we avoid self loops.

## What is a Graph?

- ▶ A graph consists of two sets  $V$  and  $E$  where
  - ▶
$$V = \{v_1, \dots, v_n\}$$
  - ▶
$$E = \{e_1, \dots, e_m\}$$
- ▶ where each edge  $e_i$  consists of an **unordered** vertex pair  $e_i = \{v_j, v_k\}$ .
- ▶  $j \neq k$  because we avoid self loops.
- ▶ For a *directed* graph we consider edges as ordered pairs.

## Example

- ▶ Let  $V = \{1, 2, 3, 4, 5\}$  and

## Example

- ▶ Let  $V = \{1, 2, 3, 4, 5\}$  and
- ▶  $E = \{(1, 2), (1, 3), (2, 3), (2, 4)\}$ .

## Example

- ▶ Let  $V = \{1, 2, 3, 4, 5\}$  and
- ▶  $E = \{(1, 2), (1, 3), (2, 3), (2, 4)\}$ .
- ▶ For an unordered graph, the order of the vertices in each edge does not matter.

## Example

- ▶ Let  $V = \{1, 2, 3, 4, 5\}$  and
- ▶  $E = \{(1, 2), (1, 3), (2, 3), (2, 4)\}$ .
- ▶ For an unordered graph, the order of the vertices in each edge does not matter.
- ▶  $G = \{V, E\}$  is a graph.